

# Newtonian gravity, red shift, confinement, asymptotic freedom and quarks oscillations

G. Quznetsov <sup>\*</sup>  
Chelyabinsk State University

October 18, 2008

## Abstract

Quarks oscillations give the Newtonian gravity law, the red shift, the confinement and the asymptotic freedom.

Let  $\langle \rho(\underline{x}), j_1(\underline{x}), j_2(\underline{x}), j_3(\underline{x}) \rangle = \langle \rho(t, \mathbf{x}), \mathbf{j}(t, \mathbf{x}) \rangle$  be a probability density 3+1-vector of any physical event [1].

Complex functions  $\varphi_1(\underline{x}), \varphi_2(\underline{x}), \varphi_3(\underline{x}), \varphi_4(\underline{x})$  exist [2] such that

$$\begin{aligned} \rho &= \sum_{s=1}^4 \varphi_s^* \varphi_s, \\ \frac{j_\alpha}{c} &= - \sum_{k=1}^4 \sum_{s=1}^4 \varphi_s^* \beta_{s,k}^{[\alpha]} \varphi_k \end{aligned} \tag{1}$$

for every such density vector (here  $\alpha \in \{1, 2, 3\}$ ,

$$\begin{aligned} \beta^{[1]} &:= \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \beta^{[2]} := \begin{bmatrix} 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{bmatrix}, \\ \beta^{[3]} &:= \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}). \end{aligned}$$

If

$$\varphi = \begin{bmatrix} \varphi_1 \\ \varphi_2 \\ \varphi_3 \\ \varphi_4 \end{bmatrix}$$

---

<sup>\*</sup>e-mails: gunn@mail.ru, quznets@yahoo.com, gunnqu@gmail.com

then [3]:

$$\frac{1}{c}\partial_t\varphi + \left(\mathbf{i}\Theta_0 + \mathbf{i}\Upsilon_0\gamma^{[5]}\right)\varphi = \begin{pmatrix} \beta^{[1]}\partial_1 + \mathbf{i}\Theta_1\beta^{[1]} + \mathbf{i}\Upsilon_1\beta^{[1]}\gamma^{[5]} + \\ +\beta^{[2]}\partial_2 + \mathbf{i}\Theta_2\beta^{[2]} + \mathbf{i}\Upsilon_2\beta^{[2]}\gamma^{[5]} + \\ +\beta^{[3]}\partial_3 + \mathbf{i}\Theta_3\beta^{[3]} + \mathbf{i}\Upsilon_3\beta^{[3]}\gamma^{[5]} + \\ +\mathbf{i}M_0\gamma_\zeta^{[0]} + \mathbf{i}M_4\beta^{[4]} - \\ -\mathbf{i}M_{\zeta,0}\gamma_\zeta^{[0]} + \mathbf{i}M_{\zeta,4}\zeta^{[4]} - \\ -\mathbf{i}M_{\eta,0}\gamma_\eta^{[0]} - \mathbf{i}M_{\eta,4}\eta^{[4]} + \\ +\mathbf{i}M_{\theta,0}\gamma_\theta^{[0]} + \mathbf{i}M_{\theta,4}\theta^{[4]} \end{pmatrix} \varphi. \quad (2)$$

with real  $\Theta_k$ ,  $\Upsilon_k$ ,  $M_0$ ,  $M_4$ ,  $M_{\zeta,0}$ ,  $M_{\zeta,4}$ ,  $M_{\eta,0}$ ,  $M_{\eta,4}$ ,  $M_{\theta,0}$ ,  $M_{\theta,4}$  and

$$\gamma^{[5]} \stackrel{def}{=} \begin{bmatrix} 1_2 & 0_2 \\ 0_2 & -1_2 \end{bmatrix}, \quad (3)$$

From [4]: *the colored motion equation* as the following:

$$\begin{pmatrix} -\beta^{[0]}\mathbf{i}\partial_0 + \Theta_0\beta^{[0]} + \Upsilon_0\beta^{[0]}\gamma^{[5]} + \\ -\beta^{[1]}\mathbf{i}\partial_1 + \Theta_1\beta^{[1]} + \Upsilon_1\beta^{[1]}\gamma^{[5]} - \\ -\beta^{[2]}\mathbf{i}\partial_2 + \Theta_2\beta^{[2]} + \Upsilon_2\beta^{[2]}\gamma^{[5]} - \\ -\beta^{[3]}\mathbf{i}\partial_3 + \Theta_3\beta^{[3]} + \Upsilon_3\beta^{[3]}\gamma^{[5]} - \\ -M_{\zeta,0}\gamma_\zeta^{[0]} + M_{\zeta,4}\zeta^{[4]} + \\ -M_{\eta,0}\gamma_\eta^{[0]} - M_{\eta,4}\eta^{[4]} + \\ +M_{\theta,0}\gamma_\theta^{[0]} + M_{\theta,4}\theta^{[4]} \end{pmatrix} \varphi = 0. \quad (4)$$

Here [5]:

$$\gamma_\zeta^{[0]} = \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{bmatrix}, \zeta^{[4]} = \begin{bmatrix} 0 & 0 & 0 & \mathbf{i} \\ 0 & 0 & \mathbf{i} & 0 \\ 0 & -\mathbf{i} & 0 & 0 \\ -\mathbf{i} & 0 & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of red pentad;

$$\gamma_\eta^{[0]} = \begin{bmatrix} 0 & 0 & 0 & \mathbf{i} \\ 0 & 0 & -\mathbf{i} & 0 \\ 0 & \mathbf{i} & 0 & 0 \\ -\mathbf{i} & 0 & 0 & 0 \end{bmatrix}, \eta^{[4]} = \begin{bmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of green pentad;

$$\gamma_\theta^{[0]} = \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix}; \theta^{[4]} = \begin{bmatrix} 0 & 0 & -\mathbf{i} & 0 \\ 0 & 0 & 0 & \mathbf{i} \\ -\mathbf{i} & 0 & 0 & 0 \\ 0 & \mathbf{i} & 0 & 0 \end{bmatrix}$$

are the antidiagonal (mass) elements of blue pentad;

$M_{\zeta,0}$ ,  $M_{\zeta,4}$  are red down and up quark's mass numbers;  
 $M_{\eta,0}$ ,  $M_{\eta,4}$  are green down and up quark's mass numbers;  
 $M_{\theta,0}$ ,  $M_{\theta,4}$  are blue down and up quark's mass numbers.

The quarks mass members of this equation form the following matrix sum:

$$\begin{aligned} \widehat{M} &\stackrel{def}{=} \begin{aligned} &-M_{\zeta,0}\gamma_{\zeta}^{[0]} + M_{\zeta,4}\zeta^{[4]} - \\ &-M_{\eta,0}\gamma_{\eta}^{[0]} - M_{\eta,4}\eta^{[4]} + \\ &+M_{\theta,0}\gamma_{\theta}^{[0]} + M_{\theta,4}\theta^{[4]} \end{aligned} \\ &= \begin{bmatrix} 0 & 0 & -M_{\theta,0} & M_{\zeta,0} - iM_{\eta,0} \\ 0 & 0 & M_{\zeta,0} + iM_{\eta,0} & M_{\theta,0} \\ -M_{\theta,0} & M_{\zeta,0} - iM_{\eta,0} & 0 & 0 \\ M_{\zeta,0} + iM_{\eta,0} & M_{\theta,0} & 0 & 0 \end{bmatrix} \\ &+ i \begin{bmatrix} 0 & 0 & -M_{\theta,4} & M_{\zeta,4} + iM_{\eta,4} \\ 0 & 0 & M_{\zeta,4} - iM_{\eta,4} & M_{\theta,4} \\ -M_{\theta,4} & -M_{\zeta,4} - iM_{\eta,4} & 0 & 0 \\ -M_{\zeta,4} + iM_{\eta,4} & M_{\theta,4} & 0 & 0 \end{bmatrix}. \end{aligned}$$

Elements of these matrices can be turned by formula of shape [6]:

$$\begin{aligned} &= \begin{pmatrix} \cos \frac{\theta}{2} & i \sin \frac{\theta}{2} \\ i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \begin{pmatrix} Z & X - iY \\ X + iY & -Z \end{pmatrix} \begin{pmatrix} \cos \frac{\theta}{2} & -i \sin \frac{\theta}{2} \\ -i \sin \frac{\theta}{2} & \cos \frac{\theta}{2} \end{pmatrix} \\ &= \begin{pmatrix} Z \cos \theta - Y \sin \theta & X - i(Y \cos \theta + Z \sin \theta) \\ X + i(Y \cos \theta + Z \sin \theta) & -Z \cos \theta + Y \sin \theta \end{pmatrix}. \end{aligned}$$

Hence, if:

$$U_{2,3}(\alpha) \stackrel{def}{=} \begin{bmatrix} \cos \alpha & i \sin \alpha & 0 & 0 \\ i \sin \alpha & \cos \alpha & 0 & 0 \\ 0 & 0 & \cos \alpha & i \sin \alpha \\ 0 & 0 & i \sin \alpha & \cos \alpha \end{bmatrix}$$

and

$$\begin{aligned} \widehat{M}' &\stackrel{def}{=} \begin{aligned} &-M'_{\zeta,0}\gamma_{\zeta}^{[0]} + M'_{\zeta,4}\zeta^{[4]} - \\ &-M'_{\eta,0}\gamma_{\eta}^{[0]} - M'_{\eta,4}\eta^{[4]} + \\ &+M'_{\theta,0}\gamma_{\theta}^{[0]} + M'_{\theta,4}\theta^{[4]} \end{aligned} \stackrel{def}{=} U_{2,3}^{\dagger}(\alpha) \widehat{M} U_{2,3}(\alpha) \end{aligned}$$

then:

$$\begin{aligned} M'_{\zeta,0} &= M_{\zeta,0}, \\ M'_{\eta,0} &= M_{\eta,0} \cos 2\alpha + M_{\theta,0} \sin 2\alpha, \\ M'_{\theta,0} &= M_{\theta,0} \cos 2\alpha - M_{\eta,0} \sin 2\alpha, \\ M'_{\zeta,4} &= M_{\zeta,4}, \\ M'_{\eta,4} &= M_{\eta,4} \cos 2\alpha + M_{\theta,4} \sin 2\alpha, \end{aligned}$$

$$M'_{\theta,4} = M_{\theta,4} \cos 2\alpha - M_{\eta,4} \sin 2\alpha.$$

Therefore, matrix  $U_{2,3}(\alpha)$  makes an oscillation between green and blue quarks colors.

Let us consider equation (2) under transformation  $U_{2,3}(\alpha)$  with  $\alpha$  is an arbitrary real function of time-space variables ( $\alpha = \alpha(t, x_1, x_2, x_3)$ ):

$$\begin{aligned} & U_{2,3}^\dagger(\alpha) \left( \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) U_{2,3}(\alpha) \varphi = \\ & = U_{2,3}^\dagger(\alpha) \left( \begin{array}{c} \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{array} \right) U_{2,3}(\alpha) \varphi. \end{aligned}$$

That is:

$$\begin{aligned} & \left( \begin{array}{c} \frac{1}{c} U_{2,3}^\dagger(\alpha) \partial_t (U_{2,3}(\alpha) \varphi) \\ + i\Theta_0 U_{2,3}^\dagger(\alpha) (U_{2,3}(\alpha) \varphi) \\ + i\Upsilon_0 U_{2,3}^\dagger(\alpha) \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) = \\ & = U_{2,3}^\dagger(\alpha) \left( \begin{array}{c} \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{array} \right) U_{2,3}(\alpha) \varphi. \end{aligned}$$

Because

$$\begin{aligned} U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) &= 1_4, \\ U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) &= \gamma^{[5]} \end{aligned}$$

then

$$\begin{aligned} & \left( \begin{array}{c} \frac{1}{c} \left( (U_{2,3}^\dagger(\alpha) \partial_t U_{2,3}(\alpha)) \varphi + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_t \varphi \right) \\ + i\Theta_0 \varphi + i\Upsilon_0 \gamma^{[5]} \varphi \end{array} \right) = \\ & = \left( \begin{array}{c} U_{2,3}^\dagger(\alpha) \beta^{[1]} \left( \begin{array}{c} (\partial_1 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_1 \varphi \\ + i\Theta_1 (U_{2,3}(\alpha) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left( \begin{array}{c} (\partial_2 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_2 \varphi \\ + i\Theta_2 (U_{2,3}(\alpha) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left( \begin{array}{c} (\partial_3 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_3 \varphi \\ + i\Theta_3 (U_{2,3}(\alpha) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{array} \right) + \\ + iM_0 U_{2,3}^\dagger(\alpha) \gamma^{[0]} (U_{2,3}(\alpha) \varphi) \\ + iM_4 U_{2,3}^\dagger(\alpha) \beta^{[4]} (U_{2,3}(\alpha) \varphi) \\ + U_{2,3}^\dagger(\alpha) \widehat{M} (U_{2,3}(\alpha) \varphi) \end{array} \right). \end{aligned}$$

Since

$$\begin{aligned} U_{2,3}^\dagger(\alpha) \gamma^{[0]} U_{2,3}(\alpha) &= \gamma^{[0]}, \\ U_{2,3}^\dagger(\alpha) \beta^{[4]} U_{2,3}(\alpha) &= \beta^{[4]} \end{aligned}$$

then

$$\begin{aligned} & \left( \frac{1}{c} \left( U_{2,3}^\dagger(\alpha) \partial_t U_{2,3}(\alpha) \right) \varphi + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_t \varphi \right) = \\ & \quad + i\Theta_0 \varphi + i\Upsilon_0 \gamma^{[5]} \varphi \\ & = \left( \begin{aligned} & U_{2,3}^\dagger(\alpha) \beta^{[1]} \left( \begin{aligned} & (\partial_1 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_1 \varphi \\ & + i\Theta_1 (U_{2,3}(\alpha) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left( \begin{aligned} & (\partial_2 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_2 \varphi \\ & + i\Theta_2 (U_{2,3}(\alpha) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left( \begin{aligned} & (\partial_3 U_{2,3}(\alpha)) \varphi + U_{2,3}(\alpha) \partial_3 \varphi \\ & + i\Theta_3 (U_{2,3}(\alpha) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{2,3}(\alpha) \varphi) \end{aligned} \right) + \\ & + iM_0 \gamma^{[0]} \varphi + iM_4 \beta^{[4]} \varphi + \widehat{M}' \varphi \end{aligned} \right) \end{aligned} \right). \end{aligned}$$

Since

$$U_{2,3}^\dagger(\alpha) \beta^{[1]} = \beta^{[1]} U_{2,3}^\dagger(\alpha)$$

then

$$\begin{aligned} & \left( U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\ & = \left( \begin{aligned} & \beta^{[1]} U_{2,3}^\dagger(\alpha) \left( \begin{aligned} & (\partial_1 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_1 \\ & + i\Theta_1 U_{2,3}(\alpha) + i\Upsilon_1 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[2]} \left( \begin{aligned} & (\partial_2 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_2 \\ & + i\Theta_2 U_{2,3}(\alpha) + i\Upsilon_2 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + U_{2,3}^\dagger(\alpha) \beta^{[3]} \left( \begin{aligned} & (\partial_3 U_{2,3}(\alpha)) + U_{2,3}(\alpha) \partial_3 \\ & + i\Theta_3 U_{2,3}(\alpha) + i\Upsilon_3 \gamma^{[5]} U_{2,3}(\alpha) \end{aligned} \right) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{aligned} \right) \varphi. \end{aligned}$$

Because

$$\begin{aligned} U_{2,3}^\dagger(\alpha) \beta^{[2]} &= (\beta^{[2]} \cos 2\alpha + \beta^{[3]} \sin 2\alpha) U_{2,3}^\dagger(\alpha), \\ U_{2,3}^\dagger(\alpha) \beta^{[3]} &= (\beta^{[3]} \cos 2\alpha - \beta^{[2]} \sin 2\alpha) U_{2,3}^\dagger(\alpha) \end{aligned}$$

then

$$\left( U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi =$$

$$= \left( \begin{aligned} & \beta^{[1]} \left( \begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_1 \\ & + i\Theta_1 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_1 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \\ & + (\beta^{[2]} \cos 2\alpha + \beta^{[3]} \sin 2\alpha) \end{aligned} \right) + \\ & \left( \begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_2 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_2 \\ & + i\Theta_2 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_2 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \\ & + (\beta^{[3]} \cos 2\alpha - \beta^{[2]} \sin 2\alpha) \end{aligned} \right) + \\ & \left( \begin{aligned} & U_{2,3}^\dagger(\alpha) \partial_3 U_{2,3}(\alpha) + U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) \partial_3 \\ & + i\Theta_3 U_{2,3}^\dagger(\alpha) U_{2,3}(\alpha) + i\Upsilon_3 U_{2,3}^\dagger(\alpha) \gamma^{[5]} U_{2,3}(\alpha) \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{aligned} \right) + \end{aligned} \right) \varphi.$$

Hence

$$\begin{aligned} & \left( \frac{1}{c} \partial_t + U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\ & = \left( \begin{aligned} & \beta^{[1]} \left( \partial_1 + U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) \\ & + \beta^{[2]} \left( \begin{aligned} & (\cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3) \\ & + U_{2,3}^\dagger(\alpha) (\cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3) U_{2,3}(\alpha) \\ & + i(\Theta_2 \cos 2\alpha - \Theta_3 \sin 2\alpha) \\ & + i(\Upsilon_2 \gamma^{[5]} \cos 2\alpha - \Upsilon_3 \gamma^{[5]} \sin 2\alpha) \end{aligned} \right) \\ & + \beta^{[3]} \left( \begin{aligned} & (\cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2) \\ & + U_{2,3}^\dagger(\alpha) (\cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2) U_{2,3}(\alpha) \\ & + i(\Theta_2 \sin 2\alpha + \Theta_3 \cos 2\alpha) \\ & + i(\Upsilon_3 \gamma^{[5]} \cos 2\alpha + \Upsilon_2 \gamma^{[5]} \sin 2\alpha) \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{aligned} \right) + \end{aligned} \right) \varphi. \quad (5) \end{aligned}$$

Let  $x'_2$  and  $x'_3$  are elements of other coordinates system such that:

$$\begin{aligned} \frac{\partial x_2}{\partial x'_2} &= \cos 2\alpha, \\ \frac{\partial x_3}{\partial x'_2} &= -\sin 2\alpha, \\ \frac{\partial x_2}{\partial x'_3} &= \sin 2\alpha, \\ \frac{\partial x_3}{\partial x'_3} &= \cos 2\alpha, \\ \frac{\partial x_0}{\partial x'_2} &= \frac{\partial x_1}{\partial x'_2} = \frac{\partial x_0}{\partial x'_3} = \frac{\partial x_1}{\partial x'_3} = 0. \end{aligned}$$

Hence:

$$\begin{aligned}
\partial'_2 \stackrel{def}{=} \frac{\partial}{\partial x'_2} &= \frac{\partial}{\partial x_0} \frac{\partial x_0}{\partial x'_2} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x'_2} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x'_2} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x'_2} \\
&= \cos 2\alpha \cdot \frac{\partial}{\partial x_2} - \sin 2\alpha \cdot \frac{\partial}{\partial x_3} \\
&= \cos 2\alpha \cdot \partial_2 - \sin 2\alpha \cdot \partial_3,
\end{aligned}$$

$$\begin{aligned}
\partial'_3 \stackrel{def}{=} \frac{\partial}{\partial x'_3} &= \frac{\partial}{\partial x_0} \frac{\partial x_0}{\partial x'_3} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x'_3} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x'_3} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x'_3} \\
&= \cos 2\alpha \cdot \frac{\partial}{\partial x_3} + \sin 2\alpha \cdot \frac{\partial}{\partial x_2} \\
&= \cos 2\alpha \cdot \partial_3 + \sin 2\alpha \cdot \partial_2.
\end{aligned}$$

Therefore from (5):

$$\begin{aligned}
&\left( \frac{1}{c} \partial_t + U_{2,3}^\dagger(\alpha) \frac{1}{c} \partial_t U_{2,3}(\alpha) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\
&= \left( \begin{aligned} &\beta^{[1]} \left( \partial_1 + U_{2,3}^\dagger(\alpha) \partial_1 U_{2,3}(\alpha) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) \\ &+ \beta^{[2]} \left( \partial'_2 + U_{2,3}^\dagger(\alpha) \partial'_2 U_{2,3}(\alpha) + i\Theta'_2 + i\Upsilon'_2 \gamma^{[5]} \right) \\ &+ \beta^{[3]} \left( \partial'_3 + U_{2,3}^\dagger(\alpha) \partial'_3 U_{2,3}(\alpha) + i\Theta'_3 + i\Upsilon'_3 \gamma^{[5]} \right) + \\ &\quad + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}' \end{aligned} \right) \varphi.
\end{aligned}$$

with

$$\begin{aligned}
\Theta'_2 &\stackrel{def}{=} \Theta_2 \cos 2\alpha - \Theta_3 \sin 2\alpha, \\
\Theta'_3 &\stackrel{def}{=} \Theta_2 \sin 2\alpha + \Theta_3 \cos 2\alpha, \\
\Upsilon'_2 &\stackrel{def}{=} \Upsilon_2 \cos 2\alpha - \Upsilon_3 \sin 2\alpha, \\
\Upsilon'_3 &\stackrel{def}{=} \Upsilon_2 \sin 2\alpha + \Upsilon_3 \cos 2\alpha.
\end{aligned}$$

Therefore, the oscillation between blue and green quarks colors curves the space of events in the  $x_2, x_3$  directions.

Similarly that: matrix

$$U_{1,3}(\vartheta) \stackrel{def}{=} \begin{bmatrix} \cos \vartheta & \sin \vartheta & 0 & 0 \\ -\sin \vartheta & \cos \vartheta & 0 & 0 \\ 0 & 0 & \cos \vartheta & \sin \vartheta \\ 0 & 0 & -\sin \vartheta & \cos \vartheta \end{bmatrix}$$

with an arbitrary real function  $\vartheta(t, x_1, x_2, x_3)$  describes the oscillation between blue and red quarks colors which curves the space of events in the  $x_1, x_3$  directions. And matrix

$$U_{1,2}(\varsigma) \stackrel{def}{=} \begin{bmatrix} \cos \varsigma - i \sin \varsigma & 0 & 0 & 0 \\ 0 & \cos \varsigma + i \sin \varsigma & 0 & 0 \\ 0 & 0 & \cos \varsigma - i \sin \varsigma & 0 \\ 0 & 0 & 0 & \cos \varsigma + i \sin \varsigma \end{bmatrix}$$

with an arbitrary real function  $\varsigma(t, x_1, x_2, x_3)$  describes the oscillation between green and red quarks colors which curves the space of events in the  $x_1, x_2$  directions.

Now, let

$$U_{0,1}(\varpi) \stackrel{def}{=} \begin{bmatrix} \cosh \varpi & -\sinh \varpi & 0 & 0 \\ -\sinh \varpi & \cosh \varpi & 0 & 0 \\ 0 & 0 & \cosh \varpi & \sinh \varpi \\ 0 & 0 & \sinh \varpi & \cosh \varpi \end{bmatrix}.$$

and

$$\widehat{M}'' \stackrel{def}{=} \begin{aligned} & -M''_{\zeta,0}\gamma_{\zeta}^{[0]} + M''_{\zeta,4}\zeta^{[4]} - \\ & -M''_{\eta,0}\gamma_{\eta}^{[0]} - M''_{\eta,4}\eta^{[4]} + \\ & + M''_{\theta,0}\gamma_{\theta}^{[0]} + M''_{\theta,4}\theta^{[4]} \end{aligned} \stackrel{def}{=} U_{0,1}^{\dagger}(\varpi) \widehat{M} U_{0,1}(\varpi)$$

then:

$$\begin{aligned} M''_{\zeta,0} &= M_{\zeta,0}, \\ M''_{\eta,0} &= (M_{\eta,0} \cosh 2\varpi - M_{\theta,4} \sinh 2\varpi), \\ M''_{\theta,0} &= M_{\theta,0} \cosh 2\varpi + M_{\eta,4} \sinh 2\varpi, \\ M''_{\zeta,4} &= M_{\zeta,4} \\ M''_{\eta,4} &= M_{\eta,4} \cosh 2\varpi + M_{\theta,0} \sinh 2\varpi, \\ M''_{\theta,4} &= M_{\theta,4} \cosh 2\varpi - M_{\eta,0} \sinh 2\varpi, \end{aligned}$$

Therefore, matrix  $U_{0,1}(\varpi)$  makes an oscillation between green and blue quarks colors with an oscillation between up and down quarks.

Let us consider equation (2) under transformation  $U_{0,1}(\varpi)$  with  $\varpi$  is an arbitrary real function of time-space variables ( $\varpi = \varpi(t, x_1, x_2, x_3)$ ):

$$\begin{aligned} & U_{0,1}^{\dagger}(\varpi) \left( \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) U_{0,1}(\varpi) \varphi = \\ & = U_{0,1}^{\dagger}(\varpi) \left( \begin{aligned} & \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ & + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ & + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M} \end{aligned} \right) U_{0,1}(\varpi) \varphi. \end{aligned}$$

Hence,

$$U_{0,1}^{\dagger}(\varpi) \left( \begin{aligned} & \frac{1}{c} \partial_t (U_{0,1}(\varpi) \varphi) \\ & + i\Theta_0 (U_{0,1}(\varpi) \varphi) + i\Upsilon_0 \gamma^{[5]} (U_{0,1}(\varpi) \varphi) \end{aligned} \right) =$$



$$= U_{0,1}^\dagger(\varpi) \left( \begin{array}{l} \beta^{[1]} \left( \begin{array}{l} \partial_1 (U_{0,1}(\varpi) \varphi) \\ +i\Theta_1 (U_{0,1}(\varpi) \varphi) + i\Upsilon_1 \gamma^{[5]} (U_{0,1}(\varpi) \varphi) \end{array} \right) + \\ +\beta^{[2]} \left( \begin{array}{l} \partial_2 (U_{0,1}(\varpi) \varphi) \\ +i\Theta_2 (U_{0,1}(\varpi) \varphi) + i\Upsilon_2 \gamma^{[5]} (U_{0,1}(\varpi) \varphi) \end{array} \right) + \\ +\beta^{[3]} \left( \begin{array}{l} \partial_3 (U_{0,1}(\varpi) \varphi) \\ +i\Theta_3 (U_{0,1}(\varpi) \varphi) + i\Upsilon_3 \gamma^{[5]} (U_{0,1}(\varpi) \varphi) \\ +iM_0 \gamma^{[0]} (U_{0,1}(\varpi) \varphi) \\ +iM_4 \beta^{[4]} (U_{0,1}(\varpi) \varphi) + \widehat{M} (U_{0,1}(\varpi) \varphi) \end{array} \right) + \end{array} \right).$$

$$U_{0,1}^\dagger(\varpi) \left( \begin{array}{l} \frac{1}{c} (\partial_t U_{0,1}(\varpi)) \varphi + U_{0,1}(\varpi) \frac{1}{c} \partial_t \varphi \\ +i\Theta_0 U_{0,1}(\varpi) \varphi + i\Upsilon_0 \gamma^{[5]} U_{0,1}(\varpi) \varphi \end{array} \right) =$$

$$= U_{0,1}^\dagger(\varpi) \left( \begin{array}{l} \beta^{[1]} \left( \begin{array}{l} (\partial_1 U_{0,1}(\varpi)) \varphi + U_{0,1}(\varpi) \partial_1 \varphi \\ +i\Theta_1 U_{0,1}(\varpi) \varphi + i\Upsilon_1 \gamma^{[5]} U_{0,1}(\varpi) \varphi \end{array} \right) + \\ +\beta^{[2]} \left( \begin{array}{l} (\partial_2 U_{0,1}(\varpi)) \varphi + U_{0,1}(\varpi) \partial_2 \varphi \\ +i\Theta_2 U_{0,1}(\varpi) \varphi + i\Upsilon_2 \gamma^{[5]} U_{0,1}(\varpi) \varphi \end{array} \right) + \\ +\beta^{[3]} \left( \begin{array}{l} (\partial_3 U_{0,1}(\varpi)) \varphi + U_{0,1}(\varpi) \partial_3 \varphi \\ +i\Theta_3 U_{0,1}(\varpi) \varphi + i\Upsilon_3 \gamma^{[5]} U_{0,1}(\varpi) \varphi \\ +iM_0 \gamma^{[0]} U_{0,1}(\varpi) \varphi \\ +iM_4 \beta^{[4]} U_{0,1}(\varpi) \varphi + \widehat{M} U_{0,1}(\varpi) \varphi \end{array} \right) + \end{array} \right).$$

Therefore:

$$\left( \begin{array}{l} U_{0,1}^\dagger(\varpi) \frac{1}{c} (\partial_t U_{0,1}(\varpi)) + U_{0,1}^\dagger(\varpi) U_{0,1}(\varpi) \frac{1}{c} \partial_t \\ +i\Theta_0 U_{0,1}^\dagger(\varpi) U_{0,1}(\varpi) + i\Upsilon_0 U_{0,1}^\dagger(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{array} \right) \varphi =$$

$$= \left( \begin{array}{l} U_{0,1}^\dagger(\varpi) \beta^{[1]} \left( \begin{array}{l} (\partial_1 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_1 \\ +i\Theta_1 U_{0,1}(\varpi) + i\Upsilon_1 \gamma^{[5]} U_{0,1}(\varpi) \end{array} \right) + \\ +U_{0,1}^\dagger(\varpi) \beta^{[2]} \left( \begin{array}{l} (\partial_2 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_2 \\ +i\Theta_2 U_{0,1}(\varpi) + i\Upsilon_2 \gamma^{[5]} U_{0,1}(\varpi) \end{array} \right) + \\ +U_{0,1}^\dagger(\varpi) \beta^{[3]} \left( \begin{array}{l} (\partial_3 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_3 \\ +i\Theta_3 U_{0,1}(\varpi) + i\Upsilon_3 \gamma^{[5]} U_{0,1}(\varpi) \\ +iM_0 U_{0,1}^\dagger(\varpi) \gamma^{[0]} U_{0,1}(\varpi) \\ +iM_4 U_{0,1}^\dagger(\varpi) \beta^{[4]} U_{0,1}(\varpi) + U_{0,1}^\dagger(\varpi) \widehat{M} U_{0,1}(\varpi) \end{array} \right) + \end{array} \right) \varphi.$$

Since:

$$\begin{aligned} U_{0,1}^\dagger(\varpi) U_{0,1}(\varpi) &= \left( \cosh 2\varpi - \beta^{[1]} \sinh 2\varpi \right), \\ U_{0,1}^\dagger(\varpi) &= \left( \cosh 2\varpi + \beta^{[1]} \sinh 2\varpi \right) U_{0,1}^{-1}(\varpi), \end{aligned}$$

$$\begin{aligned}
U_{0,1}^\dagger(\varpi) \beta^{[1]} &= \left( \beta^{[1]} \cosh 2\varpi - \sinh 2\varpi \right) U_{0,1}^{-1}(\varpi), \\
U_{0,1}^\dagger(\varpi) \beta^{[2]} &= \beta^{[2]} U_{0,1}^{-1}(\varpi), \\
U_{0,1}^\dagger(\varpi) \beta^{[3]} &= \beta^{[3]} U_{0,1}^{-1}(\varpi), \\
U_{0,1}^\dagger(\varpi) \gamma^{[0]} U_{0,1}(\varpi) &= \gamma^{[0]}, \\
U_{0,1}^\dagger(\varpi) \beta^{[4]} U_{0,1}(\varpi) &= \beta^{[4]}
\end{aligned}$$

then

$$\begin{aligned}
&\left( \begin{aligned} &(\cosh 2\varpi + \beta^{[1]} \sinh 2\varpi) U_{0,1}^{-1}(\varpi) \frac{1}{c} (\partial_t U_{0,1}(\varpi)) \\ &+ (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \frac{1}{c} \partial_t \\ &+ i\Theta_0 (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \\ &+ i\Upsilon_0 U_{0,1}^\dagger(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) \varphi = \\
&= \left( \begin{aligned} &(\beta^{[1]} \cosh 2\varpi - \sinh 2\varpi) \cdot \\ &U_{0,1}^{-1}(\varpi) \left( \begin{aligned} &(\partial_1 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_1 \\ &+ i\Theta_1 U_{0,1}(\varpi) + i\Upsilon_1 \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ \beta^{[2]} U_{0,1}^{-1}(\varpi) \left( \begin{aligned} &(\partial_2 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_2 \\ &+ i\Theta_2 U_{0,1}(\varpi) + i\Upsilon_2 \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ \beta^{[3]} U_{0,1}^{-1}(\varpi) \left( \begin{aligned} &(\partial_3 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_3 \\ &+ i\Theta_3 U_{0,1}(\varpi) + i\Upsilon_3 \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{aligned} \right) \varphi.
\end{aligned}$$

Hence:

$$\begin{aligned}
&\left( \begin{aligned} &(\cosh 2\varpi + \beta^{[1]} \sinh 2\varpi) U_{0,1}^{-1}(\varpi) \frac{1}{c} (\partial_t U_{0,1}(\varpi)) \\ &+ (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \frac{1}{c} \partial_t \\ &+ i\Theta_0 (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \\ &+ i\Upsilon_0 U_{0,1}^\dagger(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) \varphi = \\
&= \left( \begin{aligned} &(\beta^{[1]} \cosh 2\varpi - \sinh 2\varpi) \cdot \\ &U_{0,1}^{-1}(\varpi) \left( \begin{aligned} &(\partial_1 U_{0,1}(\varpi)) + U_{0,1}(\varpi) \partial_1 \\ &+ i\Theta_1 U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) + i\Upsilon_1 U_{0,1}^{-1}(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ \beta^{[2]} \left( \begin{aligned} &U_{0,1}^{-1}(\varpi) (\partial_2 U_{0,1}(\varpi)) + U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) \partial_2 \\ &+ i\Theta_2 U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) + i\Upsilon_2 U_{0,1}^{-1}(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ \beta^{[3]} \left( \begin{aligned} &U_{0,1}^{-1}(\varpi) (\partial_3 U_{0,1}(\varpi)) + U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) \partial_3 \\ &+ i\Theta_3 U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) + i\Upsilon_3 U_{0,1}^{-1}(\varpi) \gamma^{[5]} U_{0,1}(\varpi) \end{aligned} \right) + \\ &+ iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{aligned} \right) \varphi.
\end{aligned}$$

Because

$$\begin{aligned}
U_{0,1}^{-1}(\varpi) U_{0,1}(\varpi) &= 1_4, \\
U_{0,1}^{-1}(\varpi) \gamma^{[5]} U_{0,1}(\varpi) &= \gamma^{[5]}, \\
U_{0,1}^\dagger(\varpi) \gamma^{[5]} U_{0,1}(\varpi) &= \gamma^{[5]} \left( \cosh 2\varpi - \beta^{[1]} \sinh 2\varpi \right)
\end{aligned}$$

then

$$\begin{aligned}
& \left( \begin{aligned} & (\cosh 2\varpi + \beta^{[1]} \sinh 2\varpi) U_{0,1}^{-1}(\varpi) \frac{1}{c} \partial_t U_{0,1}(\varpi) \\ & + (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \frac{1}{c} \partial_t \\ & + i\Theta_0 (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \\ & + i\Upsilon_0 \gamma^{[5]} (\cosh 2\varpi - \beta^{[1]} \sinh 2\varpi) \end{aligned} \right) \varphi = \\
& = \left( \begin{aligned} & (\beta^{[1]} \cosh 2\varpi - \sinh 2\varpi) \cdot \\ & \cdot (\partial_1 + U_{0,1}^{-1}(\varpi) (\partial_1 U_{0,1}(\varpi)) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ & + \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(\varpi) (\partial_2 U_{0,1}(\varpi)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ & + \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(\varpi) (\partial_3 U_{0,1}(\varpi)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ & + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{aligned} \right) \varphi.
\end{aligned}$$

Hence:

$$\left( \begin{aligned} & \cosh 2\varpi \cdot U_{0,1}^{-1}(\varpi) \frac{1}{c} \partial_t U_{0,1}(\varpi) + \cosh 2\varpi \cdot \frac{1}{c} \partial_t + \sinh 2\varpi \cdot \partial_1 \\ & + i\Theta_0 \cosh 2\varpi + i\Upsilon_0 \gamma^{[5]} \cosh 2\varpi \\ & + \sinh 2\varpi \cdot U_{0,1}^{-1}(\varpi) (\partial_1 U_{0,1}(\varpi)) \\ & + i\Theta_1 \sinh 2\varpi \\ & + i \sinh 2\varpi \cdot \Upsilon_1 \gamma^{[5]} \\ & + \beta^{[1]} \sinh 2\varpi \cdot U_{0,1}^{-1}(\varpi) \frac{1}{c} \partial_t U_{0,1}(\varpi) - \beta^{[1]} \sinh 2\varpi \cdot \frac{1}{c} \partial_t \\ & - i\beta^{[1]} \Theta_0 \sinh 2\varpi - i\beta^{[1]} \Upsilon_0 \gamma^{[5]} \sinh 2\varpi - \beta^{[1]} \cosh 2\varpi \cdot \partial_1 \\ & - \beta^{[1]} \cosh 2\varpi \cdot U_{0,1}^{-1}(\varpi) (\partial_1 U_{0,1}(\varpi)) \\ & - i\beta^{[1]} \Theta_1 \cosh 2\varpi \\ & - i\beta^{[1]} \cosh 2\varpi \cdot \Upsilon_1 \gamma^{[5]} \\ & - \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(\varpi) (\partial_2 U_{0,1}(\varpi)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) \\ & - \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(\varpi) (\partial_3 U_{0,1}(\varpi)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) \\ & - iM_0 \gamma^{[0]} - iM_4 \beta^{[4]} - \widehat{M}'' \end{aligned} \right) \varphi = 0.$$

Therefore,

$$\left( \begin{aligned} & U_{0,1}^{-1}(\varpi) (\cosh 2\varpi \cdot \frac{1}{c} \partial_t + \sinh 2\varpi \cdot \partial_1) U_{0,1}(\varpi) \\ & + (\cosh 2\varpi \cdot \frac{1}{c} \partial_t + \sinh 2\varpi \cdot \partial_1) \\ & + i (\Theta_0 \cosh 2\varpi + \Theta_1 \sinh 2\varpi) \\ & + i (\Upsilon_0 \cosh 2\varpi + \sinh 2\varpi \cdot \Upsilon_1) \gamma^{[5]} \\ & - \beta^{[1]} \left( \begin{aligned} & U_{0,1}^{-1}(\varpi) (\cosh 2\varpi \cdot \partial_1 + \sinh 2\varpi \cdot \frac{1}{c} \partial_t) U_{0,1}(\varpi) \\ & + (\cosh 2\varpi \cdot \partial_1 + \sinh 2\varpi \cdot \frac{1}{c} \partial_t) \\ & + i (\Theta_1 \cosh 2\varpi + \Theta_0 \sinh 2\varpi) \\ & + i (\Upsilon_1 \cosh 2\varpi + \Upsilon_0 \sinh 2\varpi) \gamma^{[5]} \end{aligned} \right) \\ & - \beta^{[2]} (\partial_2 + U_{0,1}^{-1}(\varpi) (\partial_2 U_{0,1}(\varpi)) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) \\ & - \beta^{[3]} (\partial_3 + U_{0,1}^{-1}(\varpi) (\partial_3 U_{0,1}(\varpi)) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) \\ & - iM_0 \gamma^{[0]} - iM_4 \beta^{[4]} - \widehat{M}'' \end{aligned} \right) \varphi = 0 \quad (6)$$

Let  $t'$  and  $x'_1$  are elements of other coordinates system such that:

$$\begin{aligned}
\frac{\partial x_1}{\partial x'_1} &= \cosh 2\varpi, \\
\frac{\partial t}{\partial x'_1} &= \frac{1}{c} \sinh 2\varpi, \\
\frac{\partial x_1}{\partial t'} &= c \sinh 2\varpi. \\
\frac{\partial t}{\partial t'} &= \cosh 2\varpi, \\
\frac{\partial x_2}{\partial t'} &= \frac{\partial x_3}{\partial t'} = \frac{\partial x_2}{\partial x'_1} = \frac{\partial x_3}{\partial x'_1} = 0.
\end{aligned} \tag{7}$$

Hence:

$$\begin{aligned}
\partial'_t \stackrel{def}{=} \frac{\partial}{\partial t'} &= \frac{\partial}{\partial t} \frac{\partial t}{\partial t'} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial t'} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial t'} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial t'} \\
&= \cosh 2\varpi \cdot \frac{\partial}{\partial t} + c \sinh 2\varpi \cdot \frac{\partial}{\partial x_1} \\
&= \cosh 2\varpi \cdot \partial_t + c \sinh 2\varpi \cdot \partial_1,
\end{aligned}$$

That is:

$$\frac{1}{c} \partial'_t = \frac{1}{c} \cosh 2\varpi \cdot \partial_t + \sinh 2\varpi \cdot \partial_1.$$

And

$$\begin{aligned}
\partial'_1 \stackrel{def}{=} \frac{\partial}{\partial x'_1} &= \frac{\partial}{\partial t} \frac{\partial t}{\partial x'_1} + \frac{\partial}{\partial x_1} \frac{\partial x_1}{\partial x'_1} + \frac{\partial}{\partial x_2} \frac{\partial x_2}{\partial x'_1} + \frac{\partial}{\partial x_3} \frac{\partial x_3}{\partial x'_1} \\
&= \cosh 2\varpi \cdot \frac{\partial}{\partial x_1} + \sinh 2\varpi \cdot \frac{1}{c} \frac{\partial}{\partial t} \\
&= \cosh 2\varpi \cdot \partial_1 + \sinh 2\varpi \cdot \frac{1}{c} \partial_t.
\end{aligned}$$

Therefore from (6):

$$\left( \begin{array}{l} \beta^{[0]} \left( \frac{1}{c} \partial'_t + U_{0,1}^{-1}(\varpi) \frac{1}{c} \partial'_t U_{0,1}(\varpi) + i\Theta_0'' + i\Upsilon_0'' \gamma^{[5]} \right) \\ + \beta^{[1]} \left( \partial'_1 + U_{0,1}^{-1}(\varpi) \partial'_1 U_{0,1}(\varpi) + i\Theta_1'' + i\Upsilon_1'' \gamma^{[5]} \right) \\ + \beta^{[2]} \left( \partial_2 + U_{0,1}^{-1}(\varpi) \partial_2 U_{0,1}(\varpi) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]} \right) \\ + \beta^{[3]} \left( \partial_3 + U_{0,1}^{-1}(\varpi) \partial_3 U_{0,1}(\varpi) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]} \right) \\ + iM_0 \gamma^{[0]} + iM_4 \beta^{[4]} + \widehat{M}'' \end{array} \right) \varphi = 0$$

with

$$\begin{aligned}
\Theta_0'' &\stackrel{def}{=} \Theta_0 \cosh 2\varpi + \Theta_1 \sinh 2\varpi, \\
\Theta_1'' &\stackrel{def}{=} \Theta_1 \cosh 2\varpi + \Theta_0 \sinh 2\varpi, \\
\Upsilon_0'' &\stackrel{def}{=} \Upsilon_0 \cosh 2\varpi + \sinh 2\varpi \cdot \Upsilon_1, \\
\Upsilon_1'' &\stackrel{def}{=} \Upsilon_1 \cosh 2\varpi + \Upsilon_0 \sinh 2\varpi.
\end{aligned}$$

Therefore, the oscillation between blue and green quarks colors with the oscillation between up and down quarks curves the space of events in the  $t, x_1$  directions.

Similarly that: matrix

$$U_{0,2}(\phi) \stackrel{def}{=} \begin{bmatrix} \cosh \phi & i \sinh \phi & 0 & 0 \\ -i \sinh \phi & \cosh \phi & 0 & 0 \\ 0 & 0 & \cosh \phi & -i \sinh \phi \\ 0 & 0 & i \sinh \phi & \cosh \phi \end{bmatrix}$$

with an arbitrary real function  $\phi(t, x_1, x_2, x_3)$  describes the oscillation between blue and red quarks colors with the oscillation between up and down quarks which curves the space of events in the  $t, x_2$  directions. And matrix

$$U_{0,3}(\iota) \stackrel{def}{=} \begin{bmatrix} \cosh \iota + \sinh \iota & 0 & 0 & 0 \\ 0 & \cosh \iota - \sinh \iota & 0 & 0 \\ 0 & 0 & \cosh \iota - \sinh \iota & 0 \\ 0 & 0 & 0 & \cosh \iota + \sinh \iota \end{bmatrix}$$

with an arbitrary real function  $\iota(t, x_1, x_2, x_3)$  describes the oscillation between green and red quarks colors with the oscillation between up and down quarks which curves the space of events in the  $t, x_3$  directions.

From (7):

$$\begin{aligned}
\frac{\partial x_1}{\partial t'} &= c \sinh 2\varpi, \\
\frac{\partial t}{\partial t'} &= \cosh 2\varpi.
\end{aligned}$$

Because

$$\begin{aligned}
\sinh 2\varpi &= \frac{v}{\sqrt{1 - \frac{v^2}{c^2}}}, \\
\cosh 2\varpi &= \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}
\end{aligned}$$

with  $v$  as an velocity of system  $\{t', x'_1\}$  as respects to system  $\{t, x_1\}$  then




Figure 1:

$$v = \tanh 2\varpi.$$

Let

$$2\varpi := \nu(x_1) \frac{t}{x_1}$$

with

$$\nu(x_1) = \frac{\lambda}{|x_1|}, \lambda \text{ is some positive real constant.}$$

In that case

$$v(t, x_1) = \tanh \left( \nu(x_1) \frac{t}{x_1} \right)$$

and if  $g$  is an acceleration of system  $\{t', x'_1\}$  as respects to system  $\{t, x_1\}$  then

$$g(t, x_1) = \frac{\partial v}{\partial t} = \frac{\nu(x_1)}{\left( \cosh^2 \nu(x_1) \frac{t}{x_1} \right) x_1}.$$

Figure 1 shows a dependency from  $x_1$  of a system  $\{t', x'_1\}$  velocity  $v(t, x_1)$  in system  $\{t, x_1\}$ .

This velocity in point  $A$  does not equal to one in point  $B$ . Hence, an oscillator, placed in  $B$ , has got a nonzero velocity in respect to an observer, placed in




Figure 2:

point A. Therefore, by the Lorentz transformations [7], this oscillator frequency for observer, placed in point A, is less than own frequency of this oscillator (*red shift*).

Figure 2 shows a dependency from  $x_1$  of a system  $\{t', x'_1\}$  acceleration  $g(t, x_1)$  in system  $\{t, x_1\}$ .

If an immovable in system  $\{t, x_1\}$  object is placed in point K then in system  $\{t', x'_1\}$  this object must move to the left direction with acceleration  $g$  and  $g \simeq \frac{\lambda}{x_1^2}$ .

Zone from S to  $\infty$  is *the Newton Gravity Zone*.

Zone from B to C is *the Asimptotic Freedom Zone*.

And zone from C to D is *the Confinement Force Zone*.

## 1 Appendix

Let

$$\tilde{U}(\chi) \stackrel{def}{=} \begin{bmatrix} \cos \chi + i \sin \chi & 0 & 0 & 0 \\ 0 & \cos \chi + i \sin \chi & 0 & 0 \\ 0 & 0 & \cos 2\chi + i \sin 2\chi & 0 \\ 0 & 0 & 0 & \cos 2\chi + i \sin 2\chi \end{bmatrix}$$

and

$$\widehat{M}' \stackrel{def}{=} \begin{array}{c} -M'_{\zeta,0}\gamma_{\zeta}^{[0]} + M'_{\zeta,4}\zeta^{[4]} - \\ -M'_{\eta,0}\gamma_{\eta}^{[0]} - M'_{\eta,4}\eta^{[4]} + \\ + M'_{\theta,0}\gamma_{\theta}^{[0]} + M'_{\theta,4}\theta^{[4]} \end{array} \stackrel{def}{=} \widetilde{U}^{\dagger}(\chi) \widehat{M} \widetilde{U}(\chi)$$

then:

$$\begin{aligned} M'_{\zeta,0} &= (M_{\zeta,0} \cos \chi - M_{\zeta,4} \sin \chi), \\ M'_{\zeta,4} &= (M_{\zeta,4} \cos \chi + M_{\zeta,0} \sin \chi), \\ M'_{\eta,4} &= (M_{\eta,4} \cos \chi - M_{\eta,0} \sin \chi), \\ M'_{\eta,0} &= (M_{\eta,0} \cos \chi + M_{\eta,4} \sin \chi), \\ M'_{\theta,0} &= (M_{\theta,0} \cos \chi + M_{\theta,4} \sin \chi), \\ M'_{\theta,4} &= (M_{\theta,4} \cos \chi - M_{\theta,0} \sin \chi). \end{aligned}$$

Therefore, matrix  $\widetilde{U}(\chi)$  makes an oscillation between up and down quarks.

Let us consider equation (2) under transformation  $\widetilde{U}(\chi)$  with  $\chi$  is an arbitrary real function of time-space variables ( $\chi = \chi(t, x_1, x_2, x_3)$ ):

$$\begin{aligned} &\widetilde{U}^{\dagger}(\chi) \left( \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \widetilde{U}(\chi) \varphi = \\ &= \widetilde{U}^{\dagger}(\chi) \left( \begin{array}{c} \beta^{[1]} (\partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]}) + \\ + \beta^{[2]} (\partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]}) + \\ + \beta^{[3]} (\partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]}) + \\ + \widehat{M} \end{array} \right) \widetilde{U}(\chi) \varphi. \end{aligned}$$

Hence:

$$\begin{aligned} &\widetilde{U}^{\dagger}(\chi) \left( \frac{1}{c} \partial_t \left( \widetilde{U}(\chi) \varphi \right) + i\Theta_0 \widetilde{U}(\chi) \varphi + i\Upsilon_0 \gamma^{[5]} \widetilde{U}(\chi) \varphi \right) = \\ &= \widetilde{U}^{\dagger}(\chi) \left( \begin{array}{c} \beta^{[1]} \left( \partial_1 \left( \widetilde{U}(\chi) \varphi \right) + i\Theta_1 \widetilde{U}(\chi) \varphi + i\Upsilon_1 \gamma^{[5]} \widetilde{U}(\chi) \varphi \right) + \\ + \beta^{[2]} \left( \partial_2 \left( \widetilde{U}(\chi) \varphi \right) + i\Theta_2 \widetilde{U}(\chi) \varphi + i\Upsilon_2 \gamma^{[5]} \widetilde{U}(\chi) \varphi \right) + \\ + \beta^{[3]} \left( \partial_3 \left( \widetilde{U}(\chi) \varphi \right) + i\Theta_3 \widetilde{U}(\chi) \varphi + i\Upsilon_3 \gamma^{[5]} \widetilde{U}(\chi) \varphi \right) + \\ + \widehat{M} \widetilde{U}(\chi) \varphi \end{array} \right). \end{aligned}$$

Because

$$\gamma^{[5]} \widetilde{U}(\chi) = \widetilde{U}(\chi) \gamma^{[5]}$$

then



$$\begin{aligned}
& \tilde{U}^\dagger(\chi) \begin{pmatrix} \frac{1}{c} \left( \partial_t \tilde{U}(\chi) \right) \varphi + \tilde{U}(\chi) \frac{1}{c} \partial_t \varphi \\ +i\Theta_0 \tilde{U}(\chi) \varphi + \tilde{U}(\chi) i\Upsilon_0 \gamma^{[5]} \varphi \end{pmatrix} = \\
& = \tilde{U}^\dagger(\chi) \begin{pmatrix} \beta^{[1]} \begin{pmatrix} \left( \partial_1 \tilde{U}(\chi) \right) \varphi + \tilde{U}(\chi) \partial_1 \varphi \\ +i\Theta_1 \tilde{U}(\chi) \varphi + \tilde{U}(\chi) i\Upsilon_1 \gamma^{[5]} \varphi \end{pmatrix} + \\ +\beta^{[2]} \begin{pmatrix} \left( \partial_2 \tilde{U}(\chi) \right) \varphi + \tilde{U}(\chi) \partial_2 \varphi \\ +i\Theta_2 \tilde{U}(\chi) \varphi + \tilde{U}(\chi) i\Upsilon_2 \gamma^{[5]} \varphi \end{pmatrix} + \\ +\beta^{[3]} \begin{pmatrix} \left( \partial_3 \tilde{U}(\chi) \right) \varphi + \tilde{U}(\chi) \partial_3 \varphi \\ +i\Theta_3 \tilde{U}(\chi) \varphi + \tilde{U}(\chi) i\Upsilon_3 \gamma^{[5]} \varphi \end{pmatrix} + \\ +\widehat{M} \tilde{U}(\chi) \varphi \end{pmatrix}.
\end{aligned}$$

Because

$$\begin{aligned}
\beta^{[1]} \tilde{U}(\chi) &= \tilde{U}(\chi) \beta^{[1]}, \\
\beta^{[2]} \tilde{U}(\chi) &= \tilde{U}(\chi) \beta^{[2]}, \\
\beta^{[3]} \tilde{U}(\chi) &= \tilde{U}(\chi) \beta^{[3]}
\end{aligned}$$

then:

$$\begin{aligned}
& \begin{pmatrix} +\tilde{U}^\dagger(\chi) \tilde{U}(\chi) \frac{1}{c} \partial_t + \frac{1}{c} \tilde{U}^\dagger(\chi) \left( \partial_t \tilde{U}(\chi) \right) \\ +i\Theta_0 \tilde{U}^\dagger(\chi) \tilde{U}(\chi) + \tilde{U}^\dagger(\chi) \tilde{U}(\chi) i\Upsilon_0 \gamma^{[5]} \end{pmatrix} \varphi = \\
& = \begin{pmatrix} \beta^{[1]} \begin{pmatrix} \tilde{U}^\dagger(\chi) \tilde{U}(\chi) \partial_1 + \tilde{U}^\dagger(\chi) \left( \partial_1 \tilde{U}(\chi) \right) \\ +i\Theta_1 \tilde{U}^\dagger(\chi) \tilde{U}(\chi) + \tilde{U}^\dagger(\chi) \tilde{U}(\chi) i\Upsilon_1 \gamma^{[5]} \end{pmatrix} + \\ +\beta^{[2]} \begin{pmatrix} \tilde{U}^\dagger(\chi) \tilde{U}(\chi) \partial_2 + \tilde{U}^\dagger(\chi) \left( \partial_2 \tilde{U}(\chi) \right) \\ +i\Theta_2 \tilde{U}^\dagger(\chi) \tilde{U}(\chi) + \tilde{U}^\dagger(\chi) \tilde{U}(\chi) i\Upsilon_2 \gamma^{[5]} \end{pmatrix} + \\ +\beta^{[3]} \begin{pmatrix} \tilde{U}^\dagger(\chi) \tilde{U}(\chi) \partial_3 + \tilde{U}^\dagger(\chi) \left( \partial_3 \tilde{U}(\chi) \right) \\ +i\Theta_3 \tilde{U}^\dagger(\chi) \tilde{U}(\chi) + \tilde{U}^\dagger(\chi) \tilde{U}(\chi) i\Upsilon_3 \gamma^{[5]} \end{pmatrix} + \\ +\tilde{U}^\dagger(\chi) \widehat{M} \tilde{U}(\chi) \end{pmatrix} \varphi.
\end{aligned}$$

Because:

$$\tilde{U}^\dagger(\chi) \tilde{U}(\chi) = 1_4$$

then

$$\left( \frac{1}{c} \partial_t + \frac{1}{c} \tilde{U}^\dagger(\chi) \left( \partial_t \tilde{U}(\chi) \right) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi =$$

$$= \begin{pmatrix} \beta^{[1]} \left( \partial_1 + \tilde{U}^\dagger(\chi) \left( \partial_1 \tilde{U}(\chi) \right) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) + \\ + \beta^{[2]} \left( \partial_2 + \tilde{U}^\dagger(\chi) \left( \partial_2 \tilde{U}(\chi) \right) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]} \right) + \\ + \beta^{[3]} \left( \partial_3 + \tilde{U}^\dagger(\chi) \left( \partial_3 \tilde{U}(\chi) \right) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]} \right) + \\ + \tilde{U}^\dagger(\chi) \widehat{M} \tilde{U}(\chi) \end{pmatrix} \varphi.$$

Now let:

$$\widehat{U}(\kappa) \stackrel{def}{=} \begin{bmatrix} \cosh \kappa + \sinh \kappa & 0 & 0 & 0 \\ 0 & \cosh \kappa + \sinh \kappa & 0 & 0 \\ 0 & 0 & \cosh 2\kappa + \sinh 2\kappa & 0 \\ 0 & 0 & 0 & \cosh 2\kappa + \sinh 2\kappa \end{bmatrix}$$

and

$$\widehat{M}' \stackrel{def}{=} \begin{pmatrix} -M'_{\zeta,0} \gamma_\zeta^{[0]} + M'_{\zeta,4} \zeta^{[4]} - \\ -M'_{\eta,0} \gamma_\eta^{[0]} - M'_{\eta,4} \eta^{[4]} + \\ + M'_{\theta,0} \gamma_\theta^{[0]} + M'_{\theta,4} \theta^{[4]} \end{pmatrix} \stackrel{def}{=} \widehat{U}^{-1}(\kappa) \widehat{M} \widehat{U}(\kappa)$$

then:

$$\begin{aligned} M'_{\theta,0} &= (M_{\theta,0} \cosh \kappa - iM_{\theta,4} \sinh \kappa), \\ M'_{\theta,4} &= (M_{\theta,4} \cosh \kappa + iM_{\theta,0} \sinh \kappa), \\ M'_{\eta,0} &= (M_{\eta,0} \cosh \kappa - iM_{\eta,4} \sinh \kappa), \\ M'_{\eta,4} &= (M_{\eta,4} \cosh \kappa + iM_{\eta,0} \sinh \kappa), \\ M'_{\zeta,0} &= (M_{\zeta,0} \cosh \kappa + iM_{\zeta,4} \sinh \kappa), \\ M'_{\zeta,4} &= (M_{\zeta,4} \cosh \kappa - iM_{\zeta,0} \sinh \kappa). \end{aligned}$$

Therefore, matrix  $\widehat{U}(\kappa)$  makes an oscillation between up and down quarks, too.

Let us consider equation (2) under transformation  $\widehat{U}(\kappa)$  with  $\kappa$  is an arbitrary real function of time-space variables ( $\kappa = \kappa(t, x_1, x_2, x_3)$ ):

$$\begin{aligned} &\widehat{U}^{-1}(\kappa) \left( \frac{1}{c} \partial_t + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \widehat{U}(\kappa) \varphi = \\ &= \widehat{U}^{-1}(\kappa) \begin{pmatrix} \beta^{[1]} \left( \partial_1 + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) + \\ + \beta^{[2]} \left( \partial_2 + i\Theta_2 + i\Upsilon_2 \gamma^{[5]} \right) + \\ + \beta^{[3]} \left( \partial_3 + i\Theta_3 + i\Upsilon_3 \gamma^{[5]} \right) + \\ + \widehat{M} \end{pmatrix} \widehat{U}(\kappa) \varphi. \end{aligned}$$

Hence:

$$\begin{aligned} & \widehat{U}^{-1}(\kappa) \left( \frac{1}{c} \partial_t \left( \widehat{U}(\kappa) \varphi \right) + i\Theta_0 \widehat{U}(\kappa) \varphi + i\Upsilon_0 \gamma^{[5]} \widehat{U}(\kappa) \varphi \right) = \\ & = \widehat{U}^{-1}(\kappa) \left( \begin{aligned} & \beta^{[1]} \left( \partial_1 \left( \widehat{U}(\kappa) \varphi \right) + i\Theta_1 \widehat{U}(\kappa) \varphi + i\Upsilon_1 \gamma^{[5]} \widehat{U}(\kappa) \varphi \right) + \\ & + \beta^{[2]} \left( \partial_2 \left( \widehat{U}(\kappa) \varphi \right) + i\Theta_2 \widehat{U}(\kappa) \varphi + i\Upsilon_2 \gamma^{[5]} \widehat{U}(\kappa) \varphi \right) + \\ & + \beta^{[3]} \left( \partial_3 \left( \widehat{U}(\kappa) \varphi \right) + i\Theta_3 \widehat{U}(\kappa) \varphi + i\Upsilon_3 \gamma^{[5]} \widehat{U}(\kappa) \varphi \right) + \\ & + \widehat{M} \widehat{U}(\kappa) \varphi \end{aligned} \right). \end{aligned}$$

Because

$$\gamma^{[5]} \widehat{U}(\kappa) = \widehat{U}(\kappa) \gamma^{[5]}$$

then

$$\begin{aligned} & \widehat{U}^{-1}(\kappa) \left( \begin{aligned} & \left( \frac{1}{c} \partial_t \widehat{U}(\kappa) \right) \varphi + \widehat{U}(\kappa) \frac{1}{c} \partial_t \varphi \\ & + i\Theta_0 \widehat{U}(\kappa) \varphi + i\Upsilon_0 \widehat{U}(\kappa) \gamma^{[5]} \varphi \end{aligned} \right) = \\ & = \widehat{U}^{-1}(\kappa) \left( \begin{aligned} & \beta^{[1]} \left( \begin{aligned} & \left( \partial_1 \widehat{U}(\kappa) \right) \varphi + \widehat{U}(\kappa) \partial_1 \varphi \\ & + i\Theta_1 \widehat{U}(\kappa) \varphi + i\Upsilon_1 \widehat{U}(\kappa) \gamma^{[5]} \varphi \end{aligned} \right) + \\ & + \beta^{[2]} \left( \begin{aligned} & \left( \partial_2 \widehat{U}(\kappa) \right) \varphi + \widehat{U}(\kappa) \partial_2 \varphi \\ & + i\Theta_2 \widehat{U}(\kappa) \varphi + i\Upsilon_2 \widehat{U}(\kappa) \gamma^{[5]} \varphi \end{aligned} \right) + \\ & + \beta^{[3]} \left( \begin{aligned} & \left( \partial_3 \widehat{U}(\kappa) \right) \varphi + \widehat{U}(\kappa) \partial_3 \varphi \\ & + i\Theta_3 \widehat{U}(\kappa) \varphi + i\Upsilon_3 \widehat{U}(\kappa) \gamma^{[5]} \varphi \end{aligned} \right) + \\ & + \widehat{M} \widehat{U}(\kappa) \varphi \end{aligned} \right). \end{aligned}$$

Because

$$\begin{aligned} \widehat{U}^{-1}(\kappa) \beta^{[1]} &= \beta^{[1]} \widehat{U}^{-1}(\kappa), \\ \widehat{U}^{-1}(\kappa) \beta^{[2]} &= \beta^{[2]} \widehat{U}^{-1}(\kappa), \\ \widehat{U}^{-1}(\kappa) \beta^{[3]} &= \beta^{[3]} \widehat{U}^{-1}(\kappa) \end{aligned}$$

then

$$\left( \begin{aligned} & \widehat{U}^{-1}(\kappa) \left( \frac{1}{c} \partial_t \widehat{U}(\kappa) \right) + \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \frac{1}{c} \partial_t \\ & + i\Theta_0 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) + i\Upsilon_0 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \gamma^{[5]} \end{aligned} \right) \varphi =$$

$$= \begin{pmatrix} \beta^{[1]} \left( \begin{array}{l} \widehat{U}^{-1}(\kappa) \left( \partial_1 \widehat{U}(\kappa) \right) + \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \partial_1 \\ + i\Theta_1 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) + i\Upsilon_1 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \gamma^{[5]} \end{array} \right) + \\ + \beta^{[2]} \left( \begin{array}{l} \widehat{U}^{-1}(\kappa) \left( \partial_2 \widehat{U}(\kappa) \right) + \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \partial_2 \\ + i\Theta_2 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) + i\Upsilon_2 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \gamma^{[5]} \end{array} \right) + \\ + \beta^{[3]} \left( \begin{array}{l} \widehat{U}^{-1}(\kappa) \left( \partial_3 \widehat{U}(\kappa) \right) + \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \partial_3 \\ + i\Theta_3 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) + i\Upsilon_3 \widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) \gamma^{[5]} \\ + \widehat{U}^{-1}(\kappa) \widehat{M} \widehat{U}(\kappa) \end{array} \right) + \end{pmatrix} \varphi.$$

Because

$$\widehat{U}^{-1}(\kappa) \widehat{U}(\kappa) = 1_4$$

then

$$\begin{aligned} & \left( \frac{1}{c} \partial_t + \widehat{U}^{-1}(\kappa) \left( \frac{1}{c} \partial_t \widehat{U}(\kappa) \right) + i\Theta_0 + i\Upsilon_0 \gamma^{[5]} \right) \varphi = \\ & = \begin{pmatrix} \beta^{[1]} \left( \partial_1 + \widehat{U}^{-1}(\kappa) \left( \partial_1 \widehat{U}(\kappa) \right) + i\Theta_1 + i\Upsilon_1 \gamma^{[5]} \right) + \\ + \beta^{[2]} \left( \partial_2 + \widehat{U}^{-1}(\kappa) \left( \partial_2 \widehat{U}(\kappa) \right) + i\Theta_2 + i\Upsilon_2 \gamma^{[5]} \right) + \\ + \beta^{[3]} \left( \partial_3 + \widehat{U}^{-1}(\kappa) \left( \partial_3 \widehat{U}(\kappa) \right) + i\Theta_3 + i\Upsilon_3 \gamma^{[5]} \right) + \\ + \widehat{U}^{-1}(\kappa) \widehat{M} \widehat{U}(\kappa) \end{pmatrix} \varphi. \end{aligned}$$

If denote:

$$\begin{aligned} \Lambda_1 &\stackrel{def}{=} \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \Lambda_2 \stackrel{def}{=} \begin{bmatrix} 0 & i & 0 & 0 \\ i & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \end{bmatrix}, \\ \Lambda_3 &\stackrel{def}{=} \begin{bmatrix} 0 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}, \Lambda_4 \stackrel{def}{=} \begin{bmatrix} 0 & i & 0 & 0 \\ -i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i \\ 0 & 0 & i & 0 \end{bmatrix}, \\ \Lambda_5 &\stackrel{def}{=} \begin{bmatrix} -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & 0 & 0 & i \end{bmatrix}, \Lambda_6 \stackrel{def}{=} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \\ \Lambda_7 &\stackrel{def}{=} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix}, \Lambda_8 \stackrel{def}{=} \begin{bmatrix} i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & 2i & 0 \\ 0 & 0 & 0 & 2i \end{bmatrix} \end{aligned}$$

then

$$\begin{aligned}
U_{0,1}^{-1}(\varpi) (\partial_s U_{0,1}(\varpi)) &= \Lambda_1 \partial_s \varpi, \\
U_{2,3}^{-1}(\alpha) (\partial_s U_{2,3}(\alpha)) &= \Lambda_2 \partial_s \alpha, \\
U_{1,3}^{-1}(\vartheta) (\partial_s U_{1,3}(\vartheta)) &= \Lambda_3 \partial_s \vartheta, \\
U_{0,2}^{-1}(\phi) (\partial_s U_{0,2}(\phi)) &= \Lambda_4 \partial_s \phi, \\
U_{1,2}^{-1}(\varsigma) (\partial_s U_{1,2}(\varsigma)) &= \Lambda_5 \partial_s \varsigma, \\
U_{0,3}^{-1}(\iota) (\partial_s U_{0,3}(\iota)) &= \Lambda_6 \partial_s \iota, \\
\widehat{U}^{-1}(\kappa) (\partial_s \widehat{U}(\kappa)) &= \Lambda_7 \partial_s \kappa, \\
\widetilde{U}^{-1}(\chi) (\partial_s \widetilde{U}(\chi)) &= \Lambda_8 \partial_s \chi.
\end{aligned}$$

Let  $U$  is a product of elements of the following set:

$$\dot{U} \stackrel{def}{=} \left\{ U_{0,1}, U_{2,3}, U_{1,3}, U_{0,2}, U_{1,2}, U_{0,3}, \widehat{U}, \widetilde{U} \right\}.$$

For example, let  $U(\varsigma, \kappa, \iota) \stackrel{def}{=} U_{0,1}(\varpi) U_{0,2}(\phi) U_{1,3}(\vartheta)$   
In that case

$$\begin{aligned}
&U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
&(U_{0,1}(\varpi) U_{0,2}(\phi) U_{1,3}(\vartheta))^{-1} (\partial_s U_{0,1}(\varpi) U_{0,2}(\phi) U_{1,3}(\vartheta)).
\end{aligned}$$

Hence:

$$\begin{aligned}
&U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
&U_{1,3}^{-1}(\vartheta) U_{0,2}^{-1}(\phi) \Lambda_1 \partial_s \varpi U_{0,2}(\phi) U_{1,3}(\vartheta) \\
&+ U_{1,3}^{-1}(\vartheta) U_{0,2}^{-1}(\phi) (\partial_s U_{0,2}(\phi)) U_{1,3}(\vartheta) \\
&+ U_{1,3}^{-1}(\vartheta) \partial_s U_{1,3}(\vartheta).
\end{aligned}$$

Therefore:

$$\begin{aligned}
&U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
&U_{1,3}^{-1}(\vartheta) (\Lambda_1 \cosh 2\phi - \Lambda_5 \sinh 2\phi) U_{1,3}(\vartheta) \partial_s \varpi \\
&+ U_{1,3}^{-1}(\vartheta) \Lambda_4 \partial_s \phi U_{1,3}(\vartheta) \\
&+ \Lambda_3 \partial_s \vartheta.
\end{aligned}$$

Hence:

$$\begin{aligned}
&U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
&U_{1,3}^{-1}(\vartheta) \Lambda_1 \cosh 2\phi U_{1,3}(\vartheta) \partial_s \varpi \\
&- U_{1,3}^{-1}(\vartheta) \Lambda_5 \sinh 2\phi U_{1,3}(\vartheta) \partial_s \varpi \\
&+ \Lambda_4 \partial_s \phi + \Lambda_3 \partial_s \vartheta.
\end{aligned}$$

Hence:

$$\begin{aligned}
U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
(\Lambda_1 \cos 2\vartheta + \Lambda_6 \sin 2\vartheta) \cosh 2\phi \partial_s \varpi \\
- (\Lambda_5 \cos 2\vartheta - \Lambda_2 \sin 2\vartheta) \sinh 2\phi \partial_s \varpi \\
+ \Lambda_4 \partial_s \phi + \Lambda_3 \partial_s \vartheta.
\end{aligned}$$

Hence:

$$\begin{aligned}
U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
\Lambda_1 \cos 2\vartheta \cosh 2\phi \partial_s \varpi + \Lambda_6 \sin 2\vartheta \cosh 2\phi \partial_s \varpi \\
- \Lambda_5 \cos 2\vartheta \sinh 2\phi \partial_s \varpi - \Lambda_2 \sin 2\vartheta \sinh 2\phi \partial_s \varpi \\
+ \Lambda_4 \partial_s \phi + \Lambda_3 \partial_s \vartheta.
\end{aligned}$$

$$\begin{aligned}
U^{-1}(\varsigma, \kappa, \iota) (\partial_s U(\varsigma, \kappa, \iota)) = \\
\Lambda_1 \cos 2\vartheta \cosh 2\phi \partial_s \varpi - \Lambda_2 \sin 2\vartheta \sinh 2\phi \partial_s \varpi + \Lambda_3 \partial_s \vartheta \\
+ \Lambda_4 \partial_s \phi - \Lambda_5 \cos 2\vartheta \sinh 2\phi \partial_s \varpi + \Lambda_6 \sin 2\vartheta \cosh 2\phi \partial_s \varpi.
\end{aligned}$$

Similarly that, there for every product  $U$  of  $\dot{U}$ 's elements real functions  $G_s^r(t, x_1, x_2, x_3)$  exist such that

$$U^{-1}(\partial_s U) = \frac{g_3}{2} \sum_{r=1}^8 \Lambda_r G_s^r$$

with some real constant  $g_3$  (similar to 8 gluons).

---


$$\begin{aligned}
U_{2,3}^{-1}(\alpha) \Lambda_1 U_{2,3}(\alpha) &= \Lambda_1 \\
U_{1,3}^{-1}(\vartheta) \Lambda_1 U_{1,3}(\vartheta) &= (\Lambda_1 \cos 2\vartheta + \Lambda_6 \sin 2\vartheta) \\
U_{0,2}^{-1}(\phi) \Lambda_1 U_{0,2}(\phi) &= (\Lambda_1 \cosh 2\phi - \Lambda_5 \sinh 2\phi) \\
U_{1,2}^{-1}(\varsigma) \Lambda_1 U_{1,2}(\varsigma) &= \Lambda_1 \cos 2\varsigma - \Lambda_4 \sin 2\varsigma \\
U_{0,3}^{-1}(\iota) \Lambda_1 U_{0,3}(\iota) &= \Lambda_1 \cosh 2\iota + \Lambda_3 \sinh 2\iota \\
\widehat{U}^{-1}(\kappa) \Lambda_1 \widehat{U}(\kappa) &= \Lambda_1 \\
\widetilde{U}^{-1}(\chi) \Lambda_1 \widetilde{U}(\chi) &= \Lambda_1 \\
===== \\
\widetilde{U}^{-1}(\chi) \Lambda_2 \widetilde{U}(\chi) &= \Lambda_2 \\
\widehat{U}^{-1}(\kappa) \Lambda_2 \widehat{U}(\kappa) &= \Lambda_2 \\
U_{0,3}^{-1}(\iota) \Lambda_2 U_{0,3}(\iota) &= \Lambda_2 \cosh 2\iota - \Lambda_4 \sinh 2\iota \\
U_{1,2}^{-1}(\varsigma) \Lambda_2 U_{1,2}(\varsigma) &= \Lambda_2 \cos 2\varsigma - \Lambda_3 \sin 2\varsigma \\
U_{0,2}^{-1}(\phi) \Lambda_2 U_{0,2}(\phi) &= \Lambda_2 \cosh 2\phi + \Lambda_6 \sinh 2\phi \\
U_{1,3}^{-1}(\vartheta) \Lambda_2 U_{1,3}(\vartheta) &= \Lambda_2 \cos 2\vartheta + \Lambda_5 \sin 2\vartheta \\
U_{0,1}^{-1}(\varpi) \Lambda_2 U_{0,1}(\varpi) &= \Lambda_2 \\
===== \\
U_{0,1}^{-1}(\varpi) \Lambda_3 U_{0,1}(\varpi) &= \Lambda_3 \cosh 2\varpi - \Lambda_6 \sinh 2\varpi \\
U_{2,3}^{-1}(\alpha) \Lambda_3 U_{2,3}(\alpha) &= \Lambda_3 \cos 2\alpha - \Lambda_5 \sin 2\alpha
\end{aligned}$$

$$\begin{aligned}
& U_{0,2}^{-1}(\phi) \quad \Lambda_3 U_{0,2}(\phi) = \Lambda_3 \\
& U_{1,2}^{-1}(\varsigma) \quad \Lambda_3 U_{1,2}(\varsigma) = \Lambda_3 \cos 2\varsigma + \Lambda_2 \sin 2\varsigma \\
& U_{0,3}^{-1}(\iota) \quad \Lambda_3 U_{0,3}(\iota) = \Lambda_3 \cosh 2\iota + \Lambda_1 \sinh 2\iota \\
& \widehat{U}^{-1}(\kappa) \quad \Lambda_3 \widehat{U}(\kappa) = \Lambda_3 \\
& \widetilde{U}^{-1}(\chi) \quad \Lambda_3 \widetilde{U}(\chi) = \Lambda_3 \\
& ===== \\
& \widetilde{U}^{-1}(\chi) \quad \Lambda_4 \widetilde{U}(\chi) = \Lambda_4 \\
& \widehat{U}^{-1}(\kappa) \quad \Lambda_4 \widehat{U}(\kappa) = \Lambda_4 \\
& U_{0,3}^{-1}(\iota) \quad \Lambda_4 U_{0,3}(\iota) = \Lambda_4 \cosh 2\iota - \Lambda_2 \sinh 2\iota \\
& U_{1,2}^{-1}(\varsigma) \quad \Lambda_4 U_{1,2}(\varsigma) = \Lambda_4 \cos 2\varsigma + \Lambda_1 \sin 2\varsigma \\
& U_{1,3}^{-1}(\vartheta) \quad \Lambda_4 U_{1,3}(\vartheta) = \Lambda_4 \\
& U_{2,3}^{-1}(\alpha) \quad \Lambda_4 U_{2,3}(\alpha) = \Lambda_4 \cos 2\alpha - \Lambda_6 \sin 2\alpha \\
& U_{0,1}^{-1}(\varpi) \quad \Lambda_4 U_{0,1}(\varpi) = \Lambda_4 \cosh 2\varpi + \Lambda_5 \sinh 2\varpi \\
& ===== \\
& U_{0,1}^{-1}(\varpi) \quad \Lambda_5 U_{0,1}(\varpi) = \Lambda_5 \cosh 2\varpi + \Lambda_4 \sinh 2\varpi \\
& U_{2,3}^{-1}(\alpha) \quad \Lambda_5 U_{2,3}(\alpha) = \Lambda_5 \cos 2\alpha + \Lambda_3 \sin 2\alpha \\
& U_{1,3}^{-1}(\vartheta) \quad \Lambda_5 U_{1,3}(\vartheta) = (\Lambda_5 \cos 2\vartheta - \Lambda_2 \sin 2\vartheta) \\
& U_{0,2}^{-1}(\phi) \quad \Lambda_5 U_{0,2}(\phi) = \Lambda_5 \cosh 2\phi - \Lambda_1 \sinh 2\phi \\
& U_{0,3}^{-1}(\iota) \quad \Lambda_5 U_{0,3}(\iota) = \Lambda_5 \\
& \widehat{U}^{-1}(\kappa) \quad \Lambda_5 \widehat{U}(\kappa) = \Lambda_5 \\
& \widetilde{U}^{-1}(\chi) \quad \Lambda_5 \widetilde{U}(\chi) = \Lambda_5 \\
& ===== \\
& \widetilde{U}^{-1}(\chi) \quad \Lambda_6 \widetilde{U}(\chi) = \Lambda_6 \\
& \widehat{U}^{-1}(\kappa) \quad \Lambda_6 \widehat{U}(\kappa) = \Lambda_6 \\
& U_{1,2}^{-1}(\varsigma) \quad \Lambda_6 U_{1,2}(\varsigma) = \Lambda_6 \\
& U_{0,2}^{-1}(\phi) \quad \Lambda_6 U_{0,2}(\phi) = \Lambda_6 \cosh 2\phi + \Lambda_2 \sinh 2\phi \\
& U_{1,3}^{-1}(\vartheta) \quad \Lambda_6 U_{1,3}(\vartheta) = \Lambda_6 \cos 2\vartheta - \Lambda_1 \sin 2\vartheta \\
& U_{2,3}^{-1}(\alpha) \quad \Lambda_6 U_{2,3}(\alpha) = \Lambda_6 \cos 2\alpha + \Lambda_4 \sin 2\alpha \\
& U_{0,1}^{-1}(\varpi) \quad \Lambda_6 U_{0,1}(\varpi) = \Lambda_6 \cosh 2\varpi - \Lambda_3 \sinh 2\varpi \\
& ===== \\
& \widetilde{U}^{-1}(\chi) \quad \Lambda_7 \widetilde{U}(\chi) = \Lambda_7 \\
& U_{0,3}^{-1}(\iota) \quad \Lambda_7 U_{0,3}(\iota) = \Lambda_7 \\
& U_{1,2}^{-1}(\varsigma) \quad \Lambda_7 U_{1,2}(\varsigma) = \Lambda_7 \\
& U_{0,2}^{-1}(\phi) \quad \Lambda_7 U_{0,2}(\phi) = \Lambda_7 \\
& U_{1,3}^{-1}(\vartheta) \quad \Lambda_7 U_{1,3}(\vartheta) = \Lambda_7 \\
& U_{2,3}^{-1}(\alpha) \quad \Lambda_7 U_{2,3}(\varpi) = \Lambda_7 \\
& U_{0,1}^{-1}(\varpi) \quad \Lambda_7 U_{0,1}(\varpi) = \Lambda_7 \\
& ===== \\
& U_{0,1}^{-1}(\varpi) \quad \Lambda_8 U_{0,1}(\varpi) = \Lambda_8 \\
& U_{2,3}^{-1}(\alpha) \quad \Lambda_8 U_{2,3}(\alpha) = \Lambda_8 \\
& U_{1,3}^{-1}(\vartheta) \quad \Lambda_8 U_{1,3}(\vartheta) = \Lambda_8
\end{aligned}$$

$$\begin{aligned}
U_{0,2}^{-1}(\phi) \Lambda_8 U_{0,2}(\phi) &= \Lambda_8 \\
U_{1,2}^{-1}(\varsigma) \Lambda_8 U_{1,2}(\varsigma) &= \Lambda_8 \\
U_{0,3}^{-1}(\iota) \Lambda_8 U_{0,3}(\iota) &= \Lambda_8 \\
\widehat{U}^{-1}(\kappa) \Lambda_8 \widehat{U}(\kappa) &= \Lambda_8
\end{aligned}$$

## References

- [1] Gunn Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series Contemporary Fundamental Physics, ed. V. Dvoeglazov, Nova Sci. Publ. Inc., NY (2007), p. 38; G. Quznetsov, *Logical Foundation of Theoretical Physics*, Nova Sci. Publ. Inc., NY, (2006), p. 80
- [2] Gunn Quznetsov, *Probabilistic Treatment of Gauge Theories*, in series Contemporary Fundamental Physics, ed. V. Dvoeglazov, Nova Sci. Publ. Inc., NY (2007), pp.39-41
- [3] Item, p.61, eq.(2.27)
- [4] Item, p.136, eq.(3.49)
- [5] Item, p.p.38-39
- [6] For instance J. M. Ziman, *Elements of Advanced Quantum Theory*, Cambridge University Press (1969) formula (6.59)
- [7] G. Quznetsov, Logical Foundation of Theoretical Physics in series "Contemporary Fundamental Physics", Sci.Nova Publ. NY (2006) pp. 19-51